



Replacement of laparoscopic total gastrectomy with laparoscopic proximal gastrectomy for upper early gastric cancer

Susumu Shibasaki¹, Koichi Suda^{1,2}, Ichiro Uyama¹

¹Department of Surgery, Fujita Health University, 1-98 Dengakugakubo, Kutsukake, Toyoake, Aichi, Japan; ²Cancer Center, School of Medicine, Keio University, 35 Shinanomachi, Shinjyuku, Tokyo, Japan

Correspondence to: Koichi Suda, MD, PhD, FACS. Department of Surgery, Fujita Health University, 1-98 Dengakugakubo, Kutsukake, Toyoake, Aichi 470-1192, Japan. Email: ko-suda@nifty.com.

Comment on: Nishigori T, Okabe H, Tsunoda S, *et al.* Superiority of laparoscopic proximal gastrectomy with hand-sewn esophagogastronomy over total gastrectomy in improving postoperative body weight loss and quality of life. *Surg Endosc* 2017;31:3664-72.

Received: 26 January 2018; Accepted: 20 February 2018; Published: 07 March 2018.

doi: 10.21037/ales.2018.02.02

View this article at: <http://dx.doi.org/10.21037/ales.2018.02.02>

In accordance with the increased incidence of proximal early gastric cancer (EGC), particularly in Asian countries (1,2), the demand for proximal gastrectomy (PG) as a function-preserving surgery continues to escalate. Although PG offers numerous potential advantages in preserving the physiological function of the remnant stomach, a complicated reconstructive procedure is required to restructure the anti-reflux mechanism to prevent reflux esophagitis (3,4). Among the three main reconstruction methods, esophagogastronomy (EG) is performed more often than double-tract reconstruction (DTR) and jejunal interposition (JI) (3,4); this is owing to its relative simplicity, although an optimal reconstructive procedure is yet to be established. With the recent advances in laparoscopic techniques and the expanded indications for laparoscopic gastrectomy for the treatment of proximal EGC, the number of reports of novel reconstructive procedures for laparoscopic PG (LPG) with EG has been rapidly increasing over the past few years (5-11). However, these reports have focused on technical aspects and surgical outcomes rather than on long-term quality of life (QOL), which remains unclear.

Recently, Nishigori *et al.* reported the institutional standardization of laparoscopic gastrectomy even for gastric cancer (GC) located in the upper third stomach; this is because of the advantages of LPG, unlike laparoscopic total gastrectomy (LTG), in preventing postoperative weight loss and in improving QOL in patients with stage I GC (12). In a single-institution retrospective comparative study,

42 patients with stage I GC underwent LTG followed by Roux-en-Y reconstruction with functional end-to-end esophagojejunal anastomosis (13), whereas to fix the distal esophagus to the gastric anterior wall, 20 GC patients underwent LPG followed by EG with a hand-sewn esophagogastric anastomosis and fundoplication using a knifeless endoscopic linear stapler (14). Although there was no significant difference in the incidence of early complications between these two groups, anastomotic stricture more frequently occurred in the LPG group (25%) than in the LTG group (0%). Although there was no significant difference in the incidence of grade \geq B postoperative reflux gastritis based on the Los Angeles classification criteria between the two groups (5% and 2%, respectively), the percentage of weight loss at 12 months was lower in the LPG group than that in the LTG group (-10.7% *vs.* -16.3%, respectively). Furthermore, multivariate analysis revealed that LPG was associated with less body weight loss (12). Patient responses to a questionnaire survey using the Postgastrectomy Syndrome Assessment Scale (PGSAS)-45, which was distributed >1 year after operation, showed that the incidences of diarrhea and dissatisfaction with symptoms were better with LPG than those with LTG. However, this survey was limited to 17 patients who underwent TG and 11 who underwent PG (12). There were three main limitations to this study: the retrospective design, small sample size, and only a single assessment time point in the questionnaire survey. Collectively, LPG for patients with proximal

EGC offers several advantages over LTG, including less postoperative body weight loss, fewer diarrhea symptoms, and better QOL; however, anastomotic stricture occurred more frequently.

This study is of great value as the subjective symptoms were examined based on a patient-oriented survey using the PGSAS-45, which was designed to assess the severity of symptoms, living status, and QOL of the gastrectomized patients (15). Although many previous studies have mainly focused on body weight loss as an indicator of long-term nutritional status following LPG (3,4), Nishigori *et al.* successfully demonstrated that LPG is superior to LTG in terms of diarrhea and dissatisfaction with symptoms, and in maintaining body weight after operation. Overall, body weight loss after gastrectomy is induced by a reduction in gastric reservoir capacity and decrease in the number of various digestive hormones, including appetite hormone ghrelin, which lead to appetite loss, inadequate oral nutritional intake, alternation of intestinal flora, and increased peristalsis and diarrhea (3). Hence, further investigations with larger sample sizes using the study approach described by Nishigori *et al.* are warranted to clarify the association between postoperative body weight loss and QOL-related factors.

In contrast, the technical feasibility of LPG remains problematic. According to the results of a retrospective survey conducted by the Japan Society for Endoscopic Surgery, 19.7% of the patients developed LPG-associated postoperative complications with anastomotic stenosis being the most common at 6.5% (16). Particularly with LPG and EG reconstruction, anastomotic stricture is considered a major postoperative complication (range, 0–28.6%) (4–11). Nishigori *et al.* also reported that the anastomotic stricture occurred at a relatively high incidence of 25% although the rate of Clavien-Dindo grade III early complications was only 5% (12). The stricture itself directly induces dysphagia, chest discomfort, and eating disorders among other complications, resulting in deterioration in QOL. Additionally, our previous report demonstrated that anastomotic complications after laparoscopic gastrectomy, such as leakage and stricture, led to inferior long-term survival rates of patients with histologically proven T1 gastric adenocarcinoma (17). Hence, it may be necessary to overcome this high incidence of postoperative anastomotic stricture to further improve QOL and long-term survival.

Regarding the other essential issues directly related to QOL after LPG, Nishigori *et al.* reported that reflux esophagitis and grade \geq B postoperative reflux esophagitis

occurred in 5% of the patients in the LPG group (12). This result seems to be favorable as compared with a previous report on the incidence of these complications occurring at a rate of 12.5–30.8% after LPG and EG (4). This relatively low incidence may be due to restructuring of the artificial angle of His and fornix by fixation of the left side of the esophagus to the anterior gastric wall using a knifeless linear stapler (14). Conversely, grade A reflux esophagitis occurred in 20% of the patients. This finding suggests that this reconstruction method can potentially compromise QOL due to symptoms associated with reflux esophagitis. Recently, as an ideal reconstruction method to potentially reproduce the physiological anti-reflux mechanism, valvuloplastic EG (VEG) with a double flap technique (DFT) was developed (6–9). In this procedure, the distal esophagus and the site of anastomosis are implanted in the submucosal layer, and the anterior side of the anastomosis is completely covered by a seromuscular double flap. These structures and intragastric pressure generate a pressure gradient between the esophagus and stomach, substituting as a one-way valve (18). In fact, these studies demonstrated that the incidence of grade \geq B reflux esophagitis was only 0–5% (6–9), although most included only small-scale, single-center experiences. However, this procedure requires complicated suturing and ligation. Additionally, the rate of anastomotic stricture reportedly is as high as 4.7–29.1%, particularly before attaining a learning plateau (6–9). Hence, the complexity and technical difficulty of VEG-DFT are obstacles to the standardization of this reconstructive procedure after LPG. Therefore, the technical difficulty and complexity of EG must be further improved.

LPG has great potential as a radical procedure for esophagogastric junction (EGJ) cancer. According to the Japanese gastric cancer guidelines 2014, PG, and not TG, is recommended considering the optimal extent of lymphadenectomy for junctional cancer of \leq 4 cm in diameter centered within 2 cm of the EGJ (19). Several recent studies have demonstrated that low body mass index and sarcopenia were independent predictive factors of the short- and long-term outcomes of gastrectomized patients, particularly those with advanced cancer requiring multidisciplinary therapy (20–22). Therefore, avoiding TG can possibly prevent postoperative severe body weight loss of patients with advanced EGJ cancer, leading to improved QOL and survival outcomes. However, no optimal reconstructive procedure following PG for EGJ cancer has yet been established as for upper EGC. Particularly in patients with EGJ cancer, the anastomotic procedure

is often intra-mediastinally performed to secure an oral cancer-free margin. Therefore, JI or DTR is preferred considering the safety of an anastomotic procedure rather than EG, which is complicated with a high risk of reflux esophagitis and involves a larger surgical field. However, JI and DTR require a greater number of anastomoses; therefore, the anastomotic time is longer than with EG, and the long-term reservoir function of the remnant stomach after these reconstructive methods remains unclear. Therefore, the merits and demerits of EG, JI, and DTR should be cautiously considered on a case-by-case basis while selecting an optimal reconstruction method. The knowledge gained from further studies focused on the long-term function of the remnant stomach with each of these reconstructive procedures after PG would be indispensable.

Two points are particularly important to consider while selecting LPG as a replacement for LTG. First, the exact incidence of anastomotic complications in LPG must be surveyed in large-scaled prospective studies. In Japan, a nonrandomized confirmatory trial was commenced in April 2015 (Japan Clinical Oncology Group Study JCOG1401, UMIN000017155) to evaluate the safety of LTG and LPG for clinical stage I GC with the incidence of anastomotic leakage as the primary endpoints and other perioperative and long-term outcomes as the secondary endpoints (23). This study reveals the actual incidence of anastomotic complications and the safety and feasibility of LPG, although the reconstructive procedure is limited to JI or DTR. Moreover, the findings of this study are expected to help define the standards to examine the safety of EG reconstruction. Second, the application of robotic surgery can potentially relieve the technical complexity and difficulty of VEG-DFT because the use of a robotic system can facilitate precise performance in a confined surgical field with impressive dexterity through several technical properties, including stereoscopic vision, the use of wristed micro-surgical instruments and tremor filtration, and the ability to scale motion (24,25). In fact, we previously reported favorable short-term outcomes and an early learning curve with the use of robotic-assisted VEG-DFT (18). Additionally, an ideal reconstructive procedure after PG with fewer complications and a lower risk of reflux esophagitis is expected to be recognized with the full utilization of a robotic system.

In conclusion, LPG with EG reconstruction has great potential to become a remarkable, function-preserving, standard, surgical procedure for patients with upper EGC and EGJ cancer once the risks of anastomotic stricture and

reflux gastritis are resolved.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Annals of Laparoscopic and Endoscopic Surgery*. The article did not undergo external peer review.

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/ales.2018.02.02>). KS serves as an unpaid editorial board member of *Annals of Laparoscopic and Endoscopic Surgery* from Jul 2016 to Jun 2018. The other authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Ahn HS, Lee HJ, Yoo MW, et al. Changes in clinicopathological features and survival after gastrectomy for gastric cancer over a 20-year period. *Br J Surg* 2011;98:255-60.
2. Sano T, Hollowood A. Early gastric cancer: diagnosis and less invasive treatments. *Scand J Surg* 2006;95:249-55.
3. Ueda Y, Shiroshita H, Etoh T, et al. Laparoscopic proximal gastrectomy for early gastric cancer. *Surg Today* 2017;47:538-47.
4. Nakamura M, Yamaue H. Reconstruction after proximal gastrectomy for gastric cancer in the upper third of the stomach: a review of the literature published from 2000 to

2014. *Surg Today* 2016;46:517-27.
5. Yasuda A, Yasuda T, Imamoto H, et al. A newly modified esophagogastrostomy with a reliable angle of His by placing a gastric tube in the lower mediastinum in laparoscopy-assisted proximal gastrectomy. *Gastric Cancer* 2015;18:850-8.
 6. Kuroda S, Nishizaki M, Kikuchi S, et al. Double flap technique as an antireflux procedure in esophagogastrostomy after proximal gastrectomy. *J Am Coll Surg* 2016;223:e7-13.
 7. Muraoka A, Kobayashi M, Kokudo Y. Laparoscopy-assisted proximal gastrectomy with the hinged double flap method. *World J Surg* 2016;40:2419-24.
 8. Hosoda K, Yamashita K, Moriya H, et al. Laparoscopically assisted proximal gastrectomy with esophagogastrostomy using a novel "open-door" technique. *J Gastrointest Surg* 2017;21:1174-80.
 9. Hayami M, Hiki N, Nunobe S, et al. Clinical outcomes and evaluation of laparoscopic proximal gastrectomy with double-flap technique for early gastric cancer in the upper third of the stomach. *Ann Surg Oncol* 2017;24:1635-42.
 10. Yamashita Y, Yamamoto A, Tamamori Y, et al. Side overlap esophagogastrostomy to prevent reflux after proximal gastrectomy. *Gastric Cancer* 2017;20:728-35.
 11. Kosuga T, Ichikawa D, Komatsu S, et al. Feasibility and Nutritional Benefits of Laparoscopic Proximal Gastrectomy for Early Gastric Cancer in the Upper Stomach. *Ann Surg Oncol* 2015;22 Suppl 3:S929-35.
 12. Nishigori T, Okabe H, Tsunoda S, et al. Superiority of laparoscopic proximal gastrectomy with hand-sewn esophagogastrostomy over total gastrectomy in improving postoperative body weight loss and quality of life. *Surg Endosc* 2017;31:3664-72.
 13. Okabe H, Obama K, Tanaka E, et al. Intracorporeal esophagojejunal anastomosis after laparoscopic total gastrectomy for patients with gastric cancer. *Surg Endosc* 2009;23:2167-71.
 14. Okabe H, Obama K, Tanaka E, et al. Laparoscopic proximal gastrectomy with a hand-sewn esophago-gastric anastomosis using a knifeless endoscopic linear stapler. *Gastric Cancer* 2013;16:268-74.
 15. Nakada K, Ikeda M, Takahashi M, et al. Characteristics and clinical relevance of postgastrectomy syndrome assessment scale (PGSAS)-45: newly developed integrated questionnaires for assessment of living status and quality of life in postgastrectomy patients. *Gastric Cancer* 2015;18:147-58.
 16. Kitano S, Yamashita Y, Siraishi N, et al. 12th Nationwide Survey of Endoscopic Surgery in Japan (in Japanese). *J Jpn Soc Endosc Surg* 2014;19:535-40.
 17. Nagasako Y, Satoh S, Isogaki J, et al. Impact of anastomotic complications on outcome after laparoscopic gastrectomy for early gastric cancer. *Br J Surg* 2012;99:849-54.
 18. Shibasaki S, Suda K, Nakauchi M, et al. Robotic valvuloplastic esophagogastrostomy using double flap technique following proximal gastrectomy: technical aspects and short-term outcomes. *Surg Endosc* 2017;31:4283-97.
 19. Japanese Gastric Cancer Association. Japanese gastric cancer treatment guidelines 2014 (ver. 4). *Gastric Cancer* 2017;20:1-19.
 20. Fujiwara Y, Fukuda S, Tsujie M, et al. Outcome predictors for patients with stage II/III gastric cancer who undergo gastrectomy and S-1 adjuvant chemotherapy. *Oncol Lett* 2017;14:1621-7.
 21. Lee HH, Park JM, Song KY, et al. Survival impact of postoperative body mass index in gastric cancer patients undergoing gastrectomy. *Eur J Cancer* 2016;52:129-37.
 22. Zhuang CL, Huang DD, Pang WY, et al. Sarcopenia is an Independent Predictor of Severe Postoperative Complications and Long-Term Survival After Radical Gastrectomy for Gastric Cancer: Analysis from a Large-Scale Cohort. *Medicine (Baltimore)* 2016;95:e3164.
 23. Kataoka K, Katai H, Mizusawa J, et al. Non-randomized confirmatory trial of laparoscopy-assisted total gastrectomy and proximal gastrectomy with nodal dissection for clinical stage I gastric cancer: Japan clinical oncology group study JCOG1401. *J Gastric Cancer* 2016;16:93-7.
 24. Suda K, Man IM, Ishida Y, et al. Potential advantages of robotic radical gastrectomy for gastric adenocarcinoma in comparison with conventional laparoscopic approach: a single institutional retrospective comparative cohort study. *Surg Endosc* 2015;29:673-85.
 25. Uyama I, Kanaya S, Ishida Y, et al. Novel integrated robotic approach for suprapancreatic D2 nodal dissection for treating gastric cancer: technique and initial experience. *World J Surg* 2012;36:331-7.

doi: 10.21037/ales.2018.02.02

Cite this article as: Shibasaki S, Suda K, Uyama I. Replacement of laparoscopic total gastrectomy with laparoscopic proximal gastrectomy for upper early gastric cancer. *Ann Laparosc Endosc Surg* 2018;3:17.